

TexaClear® Pain Reliever

Drug Facts

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product package labeling, this website labeling should be regarded as the most current.

Drug Facts	
Active ingredients (in each 30 mL)	Purpose
Caffeine 65 mg	Pain reliever aid
Choline Salicylate (NSAID*) 870 mg.....	Pain reliever / fever reducer
*nonsteroidal anti-inflammatory drug	
Uses	
temporarily relieves minor aches and pains due to:	
<ul style="list-style-type: none"> • headache • minor arthritis pain • muscular aches • menstrual cramps • backache • temporarily reduces fever 	
Warnings	
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.	
Allergy alert: Salicylates (NSAIDs) may cause a severe allergic reaction which may include: • hives • skin reddening • rash • facial swelling • shock • asthma (wheezing)	
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:	
<ul style="list-style-type: none"> • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed 	
Do not use if you have ever had an allergic reaction to salicylates (including aspirin) or any other pain reliever / fever reducer.	
Ask a doctor before use if	
<ul style="list-style-type: none"> • stomach bleeding warning applies to you • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have a history of stomach problems such as heartburn, upset stomach, stomach pain, or ulcers • you have asthma 	
Ask a doctor or pharmacist before use if you are	
<ul style="list-style-type: none"> • under a doctor's care for any serious condition • taking a prescription drug for diabetes, gout, or arthritis 	
When using this product	
<ul style="list-style-type: none"> • limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat • the recommended dose of this product contains about as much caffeine as a cup of coffee • take with food or milk if stomach upset occurs 	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • symptoms do not improve • pain gets worse or lasts more than 10 days • ringing in the ears or loss of hearing occurs • redness or swelling is present 	

Drug Facts (Continued)
<ul style="list-style-type: none"> • any new symptoms appear • fever gets worse or lasts more than 3 days
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use salicylates (NSAIDs) during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
<ul style="list-style-type: none"> • do not take more than directed • adults and children 12 years of age and over: take 30 mL every 6 hours, while symptoms persist. • do not exceed 4 doses in 24 hours unless directed by a doctor. • children under 12 years of age: ask a doctor. • use enclosed dose cup.
Other information
<ul style="list-style-type: none"> • each 30mL contains: Sodium 15 mg • read all product information before using • store at room temperature 68-86°F (20-30°C) • avoid excessive heat and humidity • tamper evident: do not use if foil seal under cap is broken or missing.
Inactive ingredients
citric acid anhydrous, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol
Questions or comments?
Call 1-888-535-0305 9 a.m. - 5 p.m. CST.



Distributed by:
GM Pharmaceuticals, Inc.
Arlington, TX. 76015

TexaClear® Pain Reliever One Dose

Drug Facts

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product package labeling, this website labeling should be regarded as the most current.

Drug Facts	
Active ingredients (per 57 mL)	Purpose
Caffeine 65 mg	Pain reliever aid
Choline Salicylate (NSAID*) 870 mg.....	Pain reliever / fever reducer
*nonsteroidal anti-inflammatory drug	
Uses	
temporarily relieves minor aches and pains due to:	
<ul style="list-style-type: none"> • headache • minor arthritis pain • muscular aches • menstrual cramps • backache • temporarily reduces fever 	
Warnings	
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.	
Allergy alert: Salicylates (NSAIDs) may cause a severe allergic reaction which may include: • hives • skin reddening • rash • facial swelling • shock • asthma (wheezing)	
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:	
<ul style="list-style-type: none"> • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed 	
Do not use if you have ever had an allergic reaction to salicylates (including aspirin) or any other pain reliever / fever reducer.	
Ask a doctor before use if	
<ul style="list-style-type: none"> • stomach bleeding warning applies to you • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have a history of stomach problems such as heartburn, upset stomach, stomach pain, or ulcers • you have asthma 	
Ask a doctor or pharmacist before use if you are	
<ul style="list-style-type: none"> • under a doctor's care for any serious condition • taking a prescription drug for diabetes, gout, or arthritis 	
When using this product	
<ul style="list-style-type: none"> • limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat • the recommended dose of this product contains about as much caffeine as a cup of coffee • take with food or milk if stomach upset occurs 	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • symptoms do not improve • pain gets worse or lasts more than 10 days • ringing in the ears or loss of hearing occurs • redness or swelling is present 	

Drug Facts (Continued)
<ul style="list-style-type: none"> • any new symptoms appear • fever gets worse or lasts more than 3 days
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use salicylates (NSAIDs) during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
<ul style="list-style-type: none"> • do not take more than directed • adults and children 12 years of age and over: take 1 (57 mL) bottle every 6 hours, while symptoms persist. • do not take more than 4 (57 mL) bottles in 24 hours unless directed by a doctor. • children under 12 years of age: ask a doctor.
Other information
<ul style="list-style-type: none"> • each 57 mL contains: Sodium 30 mg • read all product information before using • store at room temperature 68-86°F (20-30°C) • avoid excessive heat and humidity • tamper evident: do not use if neckband on bottle cap is broken or missing
Inactive ingredients
citric acid anhydrous, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol
Questions or comments?
Call 1-888-535-0305 9 a.m. - 5 p.m. CST.



Distributed by:
GM Pharmaceuticals, Inc.
Arlington, TX. 76015